

1 Optimization problems from high school

Let us start with the following high school math exercise question. A pharmaceutical factory manufactures two products X and Y . To produce them, we need two kinds of base materials A and B . The following table shows the amount of each material required to produce one unit of each product. Assume that we have 150 units of material A and 70 units of material B . If our objective

| | A | B |
|-----|-----|-----|
| X | 2 | 2 |
| Y | 3 | 1 |

Table 1: Materials required to produce products

is to produce as many products as possible regardless of X and Y , what is the maximum number of products we can produce with 150 units of A and 70 units of B ?

How do we solve the problem? With some abuse of notation, let x be the number of product X and y be the number of product Y that we produce. As products x and y require 2 and 3 units of material A , respectively, $2x + 3y$ would be the total amount of material A used. Here, as we have 150 units of material A , we have the condition that

$$2x + 3y \leq 150.$$

We can draw the set of points (x, y) satisfying the condition in the 2-dimensional coordinate system, and we obtain the first figure in Figure 1.1.




Figure 1.1: Deriving the set of possible solutions

Likewise, considering material B , we deduce the condition that

$$2x + y \leq 70.$$

The corresponding area in the xy -coordinate system is depicted as in the second figure of Figure 1.1. Here, of course, the quantities x and y should be nonnegative, so we have the condition that

$$x, y \geq 0.$$

In fact, a possible production plan for X and Y , given by the quantities (x, y) , must satisfy all these three conditions. The corresponding region is given as in the third figure of Figure 1.1. By simple algebra, we may deduce that the red and blue line segments intersect at point $(15, 40)$.

We have just characterized all the possible production plans under the current inventory of materials. Among these production plans, what would be the one maximizing the total number of products produced? We can simply choose the point (x, y) maximizing $x + y$. We can obtain the point by drawing line segments parallel to $x + y = 0$. We move the line $x + y$ to the upper-right corner until it does not intersect the green area. The final line is given by $x + y = 55$ going through the point $(15, 40)$ (See Figure 1.2).




Figure 1.2: Production plan maximizing the total number of produced units

Next we consider another math problem from high school. Consider a quadratic function $f(x) = 3x^2 + 6x + 5$. At which point x the function $f(x)$ is minimized? We can answer the question by




Figure 1.3: Quadratic minimization

taking the derivative of f , that is, $f'(x) = 6x + 6$. Since the coefficient of the quadratic term is positive, we know that f is minimized precisely when $f'(x) = 0$ and $x = -1$.

2 More concrete optimization problems

As you have already guessed, the two problems from high school are some examples of **optimization** problems. Before we formally define what an optimization problem is, let us discuss a few more concrete examples.

2.1 Sports game scheduling

Major League Baseball (MLB) consists of the National League (NL) and the American League (AL), each of which has 15 teams, so there are 30 teams in total. There are 162 games for all teams over 6 months in 2023, which amounts to 2430 games for one season. For example, Figure 1.4 shows the 2023 season schedule of Pittsburgh Pirates.

There are many factors and requirements when scheduling matches over the season.



P 2023 SCHEDULE

MARCH & APRIL

| SUNDAY | MONDAY | TUESDAY | WEDNESDAY | THURSDAY | FRIDAY | SATURDAY |
|----------------|----------------|----------------|-----------------|----------------|----------------|----------------|
| | | | | CIN 30 4:10 | | CIN 1 4:10 |
| CIN 2 1:40 | BOS 3 7:10 | BOS 4 7:10 | BOS 5 1:05 | | SD 6 4:10 | SD 8 8:05 |
| SD 9 1:05 | STL 10 6:55 | STL 11 6:55 | STL 12 12:05 | STL 13 7:45 | STL 14 8:15 | STL 15 2:15 |
| STL 16 2:15 | COL 17 8:40 | COL 18 8:40 | COL 19 3:10 | C 20 8:35 | C 21 8:35 | C 22 8:35 |
| C 23 1:05 | | LA 24 8:35 | LA 25 8:35 | LA 26 7:05 | WSH 27 7:05 | WSH 28 1:05 |
| WSH 30 1:05 | | | | | | |

MAY

| SUNDAY | MONDAY | TUESDAY | WEDNESDAY | THURSDAY | FRIDAY | SATURDAY |
|----------------|---------------|----------------|----------------|----------------|-----------------|----------------|
| | | TB 2 8:40 | TB 3 8:40 | TB 4 1:10 | | SD 8 8:05 |
| SD 7 1:05 | SD 9 6:55 | SD 10 6:55 | SD 11 12:05 | BAL 12 7:05 | BAL 13 7:05 | |
| BAL 14 1:05 | | DET 15 7:10 | DET 16 1:10 | | A 18 4:05 | A 20 4:05 |
| A 21 1:05 | T 22 8:35 | T 23 8:35 | T 24 12:05 | | SEA 25 10:10 | SEA 27 4:10 |
| SEA 28 4:10 | SF 29 8:45 | SF 30 8:45 | SF 31 3:45 | | | |

JUNE

| SUNDAY | MONDAY | TUESDAY | WEDNESDAY | THURSDAY | FRIDAY | SATURDAY |
|----------------|--------------|----------------|----------------|----------------|----------------|----------------|
| | | | | | SD 1 7:05 | SD 2 4:05 |
| SD 4 1:05 | As 5 7:05 | As 6 7:05 | As 7 12:05 | | N 16 4:10 | N 17 4:10 |
| N 11 7:05 | | CHC 13 8:05 | CHC 14 8:05 | CHC 15 8:05 | MIL 16 8:10 | MIL 17 4:10 |
| MIL 18 2:10 | C 19 7:05 | C 20 7:05 | C 21 12:05 | MA 22 8:40 | MA 23 8:40 | MA 24 4:10 |
| MA 25 1:40 | | SD 26 7:05 | SD 27 7:05 | SD 28 12:05 | SD 29 7:05 | SD 30 7:05 |

JULY

| SUNDAY | MONDAY | TUESDAY | WEDNESDAY | THURSDAY | FRIDAY | SATURDAY |
|----------------|----------------|---------------|----------------|----------------|-----------------|----------------|
| | | | | | | ARI 1 8:05 |
| ARI 2 1:05 | LAD 3 10:10 | LAD 4 8:10 | LAD 5 10:10 | LAD 6 10:10 | ARI 7 8:40 | ARI 8 4:10 |
| ARI 9 4:10 | ALL STAR BREAK | | | | SD 14 7:05 | SD 15 7:05 |
| SD 16 12:05 | C 17 7:05 | C 18 7:05 | C 19 12:05 | | LAA 21 12:05 | LAA 22 9:07 |
| LAA 23 4:07 | SD 24 8:40 | SD 25 8:40 | SD 26 4:10 | | P 28 7:05 | P 29 7:05 |
| P 30 1:05 | | | | | | |

AUGUST

| SUNDAY | MONDAY | TUESDAY | WEDNESDAY | THURSDAY | FRIDAY | SATURDAY |
|----------------|----------------|----------------|----------------|---------------|----------------|----------------|
| | | D 1 7:05 | D 2 12:05 | MIL 3 7:15 | MIL 4 8:10 | MIL 5 7:10 |
| MIL 6 2:10 | A 7 7:05 | A 8 7:05 | A 9 7:05 | A 10 1:05 | C 11 7:05 | C 12 7:05 |
| C 13 1:05 | NYM 14 7:10 | NYM 15 1:10 | NYM 16 1:10 | | MIN 17 8:10 | MIN 18 7:10 |
| MIN 20 2:10 | SD 21 7:05 | SD 22 7:05 | SD 23 12:05 | C 24 7:05 | C 25 7:10 | C 26 7:10 |
| C 27 1:05 | KC 28 8:10 | KC 29 8:10 | KC 30 8:10 | | | |

SEPTEMBER & OCTOBER

| SUNDAY | MONDAY | TUESDAY | WEDNESDAY | THURSDAY | FRIDAY | SATURDAY |
|----------------|---------------|----------------|----------------|----------------|----------------|----------------|
| | | | | | STL 1 8:10 | STL 2 7:10 |
| STL 3 2:10 | SD 4 12:05 | SD 5 8:10 | SD 6 12:05 | | ATL 7 7:30 | ATL 8 7:30 |
| ATL 10 1:30 | W 11 8:10 | W 12 8:10 | W 13 8:10 | W 14 12:05 | NY 15 7:30 | NY 16 7:30 |
| NY 17 1:30 | | CHC 19 7:40 | CHC 20 7:40 | CHC 21 7:40 | CIN 22 8:40 | CIN 23 8:40 |
| CIN 24 1:40 | | PHI 25 8:40 | PHI 27 8:40 | PHI 28 1:00 | M 29 8:05 | M 30 8:05 |
| M 31 3:05 | | | | | | |

Figure 1.4: Pittsburgh Pirates game schedule 2023

- The number of home games and that of away games over the season should be balanced.
- The number of home games and that of away games over the weekends in a month should be balanced.
- Games are placed in series: 3–4 consecutive games with the same team.
- 2–3 consecutive series for a home stand.
- Moderate traveling for away games.
- Requirements for interleague games (NL vs AL) and inter-division games (NL East vs NL West).
- Lots of other requirements: rival matches (Yankees vs Mets, Cubs vs White Sox).

Let alone the number of games that we have to schedule, these requirements and restrictions placed on teams are what make the scheduling problem extremely difficult. At the same time, coming up with right schedules is crucial in terms of profits. MLB is indeed a huge business recording \$11 billion revenue in 2022. Attendance and league fairness heavily depend on a schedule.

In fact, finding a feasible schedule is not so trivial. To see this, let us play with a toy example.

1. There are 6 teams (A,B,C,D,E,F), and all teams play each other once.
2. There 5 days, and we schedule 3 games per day.
3. Each team plays exactly once a day.

The following table is a partial schedule of the league. Let us complete it.

| | Game 1 | Game 2 | Game 3 |
|-------|--------|--------|--------|
| Day 1 | | (C,D) | (E,F) |
| Day 2 | (A,C) | | (D,F) |
| Day 3 | (A,D) | (B,F) | (C,E) |
| Day 4 | (A,E) | | |
| Day 5 | | | |

Table 2: Partial schedule of games

First, Game 1 for Day 1 has to be (A,B), and Game 2 for Day 2 should be (B,E). Next, Game 2 on Day 4 can be (B,C), which forces Game 2 on Day 4 to be (D,F). However, teams D and F have already played on Day 2, a conflict. Therefore, we'd better have (B,D) for Day 4. Then we may complete the rest. This example demonstrates that for sports scheduling problems, finding a

| | Game 1 | Game 2 | Game 3 |
|-------|--------|--------------|--------|
| Day 1 | (A,B) | (C,D) | (E,F) |
| Day 2 | (A,C) | (B,E) | (D,F) |
| Day 3 | (A,D) | (B,F) | (C,E) |
| Day 4 | (A,E) | (B,D) | (C,F) |
| Day 5 | (A,F) | (B,C) | (D,E) |

Table 3: Complete schedule

single schedule that satisfies all the requirements is not easy. (Note: the contents and materials in this section were directly or indirectly taken from Michael Trick's talk on sports scheduling given at Bucknell University [1].)

2.2 Least squares

Next we consider a linear regression problem in statistics. For simplicity, let us assume that there is a single feature a as well as the response b . Our hypothesis is that there exists some x such that

$$b = ax,$$

which means that the response b is simply the product of the feature a and the number x . We infer the value of x based on the data of pair (a, b) as in Figure 1.5.




Figure 1.5: Linear regression

Suppose that we have n data points

$$(a_1, b_1), \dots, (a_n, b_n).$$

The method of least squares proposes a value for x as the one minimizing the following expression.

$$\frac{1}{n} \sum_{i=1}^n (b_i - a_i x)^2.$$

For this optimization problem, any number $x \in \mathbb{R}$ can be a valid solution, in contrast to the previous example of sports scheduling. In fact, taking the derivative of the function with respect to x , we obtain that

$$x = \frac{\sum_{i=1}^n a_i b_i}{\sum_{i=1}^n a_i^2}$$

minimizes the quadratic function. However, for regression, we often deal with millions of data points, in which case the number n is huge. Hence, the focus here is to develop an algorithm that iteratively finds good solutions fast.

3 What we cover and focus on throughout the course

We have discussed the production planning problem with inequalities, the sports scheduling problem, the problem of finding the minimum of a parabola, and the linear regression problem. The first two problems involve nontrivial constraints, coming up with a solution satisfying which is not always easy. This is the case in many general classes of optimization problems that arise in operations research. For example,

- Linear programming,
- Integer programming,
- Stochastic programming,
- Network flow models.

The main objective in these problem classes is to find a solution satisfying possibly complex functional constraints while optimizing the given objective function.

On the other hand, we face the quadratic minimization problems in applications with large data sets, although they are typically unconstrained or the constraint structure is easy to deal with. As

coming up with a solution is relatively easy, many solution methods for solving such problems are iterative, in that we keep updating solutions until we reach a certain stopping criteria. We learn these topics in IE539 Convex Optimization, but NOT in this course. Instead, we focus on the first kind of optimization problems.

References

- [1] Michael Trick, Sports Scheduling: Optimization and Analytics, Analytics and Operations Management Seminar at Bucknell University (October 28, 2021) [2.1](#)