

1 Outline

In this lecture, we consider

- Operations preserving convexity,
- Introduction to convex optimization,
- Applications (Portfolio optimization, Uncertainty quantification, Support vector machine)

2 Operations preserving convexity

For many problems, it is important to recognize underlying convex structures. We can determine whether certain sets and functions are convex by understanding basic rules. Moreover, based on these rules, we can build complex convex sets and functions from simpler ones.

2.1 Set operations

We first consider set operations that preserve convexity.

- Intersection: The intersection of any (possibly infinite) collection of convex sets is convex.
- Scaling: Given a convex set C and $\alpha \in \mathbb{R}$,

$$\alpha C = \{\alpha x : x \in C\}.$$

- Minkowski sum: Given convex sets $C_i \subseteq \mathbb{R}^d$ for $i = 1, \dots, k$, the Minkowski sum of them, defined by

$$C_1 + \dots + C_k = \{x^1 + \dots + x^k : x^i \in C_i \text{ for } i = 1, \dots, k\}$$

is convex.

- Cartesian Product: Given convex sets $C_i \subseteq \mathbb{R}^{d_i}$ for $i = 1, \dots, k$, the Cartesian product of them, defined by

$$C_1 \times \dots \times C_k = \{(x^1, \dots, x^k) \in \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_k} : x^i \in C_i \text{ for } i = 1, \dots, k\}$$

is convex.

- Affine image: Given a convex set C and matrices $A \in \mathbb{R}^{p \times d}$, $b \in \mathbb{R}^p$, we define an affine mapping $f(x) = Ax + b : \mathbb{R}^d \rightarrow \mathbb{R}^p$. Then

$$f(C) = \{Ax + b : x \in C\}.$$

- Inverse affine image: Given a convex set C and matrices $A \in \mathbb{R}^{p \times d}$, $b \in \mathbb{R}^p$, we define an affine mapping $f(x) = Ax + b : \mathbb{R}^d \rightarrow \mathbb{R}^p$. Then

$$f^{-1}(C) = \{x : Ax + b \in C\}.$$

2.2 Function operations

We next consider function operations preserving convexity.

- Nonnegative weighted sum: Let $f_1, \dots, f_k : \mathbb{R}^d \rightarrow \mathbb{R}$ be convex functions. Then for any $\alpha_1, \dots, \alpha_k \geq 0$,

$$\alpha_1 f_1 + \dots + \alpha_k f_k$$

is convex.

- Maximum of arbitrary collection of convex functions: Let $\{f_\gamma\}_{\gamma \in \Gamma}$ be a collection of convex functions. Then $\max_{\gamma \in \Gamma} f_\gamma$ is also convex. Here, Γ may be infinite.
- Minimizing out variables: Let $g(x, y)$ be convex function in (x, y) . Define f by $f(x) = \inf_{y \in C} g(x, y)$ for some convex set C . Then f is convex.
- Perspective function: Let $g(x)$ be a convex function. Then $f(x, t) = tg(x/t)$ is a convex function in $(x, t) \in \mathbb{R}^d \times \mathbb{R}_{++}$. Here, f is called the perspective of g .
- Affine composition: Let $g : \mathbb{R}^p \rightarrow \mathbb{R}$ be a convex function, and take matrices $A \in \mathbb{R}^{p \times d}$, $b \in \mathbb{R}^p$. Then $f : \mathbb{R}^d \rightarrow \mathbb{R}$ defined by $f(x) = g(Ax + b)$ is convex.
- Compositions: Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be a univariate non-decreasing convex function, and let $g : \mathbb{R}^d \rightarrow \mathbb{R}$ be convex. Then $f = h \circ g$ is convex.

Example 4.1. Let C be an arbitrary set of locations. Note that

$$f_1(x) = \max_{y \in C} \|x - y\|$$

measures the longest distance from x to a location in C , and

$$f_2(x) = \min_{y \in C} \|x - y\|$$

measures the shortest distance from x to a location in C . Let us show that both f_1 and f_2 are convex. Observe first that

$$g(x, y) = \|x - y\|$$

is convex in x and y . Then f_1 is convex as it is the pointwise maximum of some convex functions. Furthermore, if C is convex, then f_2 is convex because it is a partial minimization of a convex function. In summary, f_1 is convex regardless of whether C is convex or not, while f_2 is convex if the set C is convex.

3 Convex optimization problem

3.1 Basic optimization terminologies

Given a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and a set $C \subseteq \mathbb{R}^d$, we want to solve

$$\begin{aligned} & \text{minimize} && f(x) \\ & \text{subject to} && x \in C. \end{aligned} \tag{P}$$

Terminology 1:

- x is called the *decision vector*, and the components of x are called the *decision variables*. For example, decision variables capture how much to invest for a financial portfolio or where to build a hospital in a village.
- f is called the *objective function* or *cost function*. For example, the cost of a production plan.
- C is called the *domain*, *feasible region*, or *constraint set*. For example, production capacities, budget constraints.

Terminology 2:

- Any vector $x \in C$ is called a *feasible solution*
- We say that (P) is *feasible* if $C \neq \emptyset$. Otherwise, (P) is *infeasible*.
- If there exists $x \in C$ such that $f(x) \leq r$ for any $r \in \mathbb{R}$, then (P) is *unbounded*.
- If there exists some $r \in \mathbb{R}$ such that $f(x) \geq r$ for all $x \in C$, then (P) is *bounded*.

Example 4.2. When $C = \{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 \leq 1, x_1 + x_2 \geq 2\}$, then the problem is infeasible. When $f(x) = (x - 2)^2$ and $C = [-1, 5]$, then the problem is feasible and bounded.

Terminology 3:

- $\text{OPT} := \min_{x \in C} f(x)$ is the *optimal value* of the optimization problem. Then

$$\text{OPT} = \begin{cases} +\infty, & \text{if infeasible,} \\ -\infty, & \text{if feasible but unbounded,} \\ \text{finite,} & \text{if feasible and bounded.} \end{cases}$$

- A solution $x^* \in C$ such that $f(x^*) = \text{OPT}$ is called an *optimal solution*.
- We say that (P) is *solvable* if an optimal solution exists. If not, (P) is *unsolvable*.

Example 4.3. When $f(x) = (x - 2)^2$ and $C = (3, 5]$, the problem is feasible and bounded but unsolvable.

3.2 Convex optimization

When the objective function f is convex and the feasible region is a convex set, then the optimization problem (P) is referred to as a *convex optimization* or *convex minimization* problem. By using the indicator function for C , we can rewrite (P) as

$$\begin{aligned} & \text{minimize} && f(x) \\ & \text{subject to} && I_C(x) \leq 0 \end{aligned}$$

or

$$\text{minimize} \quad f(x) + I_C(x).$$

Here, f , I_C , and $f + I_C$ are all convex. The standard form of a convex optimization problem is

$$\begin{aligned} & \text{minimize} && f(x) \\ & \text{subject to} && g_i(x) \leq 0, \quad i = 1, \dots, p, \\ & && h_i(x) = 0, \quad i = 1, \dots, q \end{aligned} \tag{P'}$$

where

- the objective function f is convex,
- the inequality constraint functions g_1, \dots, g_p are convex, and
- the equality constraint functions h_1, \dots, h_q are affine.

Exercise 4.4. If g_1, \dots, g_p are convex and h_1, \dots, h_q are affine functions, $C := \{x \in \mathbb{R}^d : g_i(x) \leq 0 \text{ for } i = 1, \dots, p, h_j(x) = 0 \text{ for } j = 1, \dots, q\}$ is a convex set.

Note that

$$\min_{x \in C} f(x) = (-1) \times \max_{x \in C} -f(x)$$

and $-f$ is concave when f is convex. Therefore, the problem of maximizing a concave function over a convex domain is also a convex optimization problem.

4 Convex optimization applications

4.1 Portfolio optimization

Given d financial assets (stocks, bonds, etc), we want to allocate x_i fraction of our budget to asset $i \in [d]$. Hence, we impose condition

$$1^\top x = 1.$$

Here, $x_i < 0$ indicates a *short position*, which means borrowing shares, selling now, and returning the shares later, while $x_i \geq 0$ indicates a *long position*, buying shares now. Then $\|x\|_1 = \sum_{i=1}^d |x_i|$ means *leverage*.

Let p_i be the initial price of asset i , and p'_i be its price at the end of one period. Then the return of asset i can be defined as

$$r_i := (p'_i - p_i)/p_i.$$

Moreover, the return of my portfolio can be measured by $r^\top x$. Here, r is a random variable with mean μ and covariance Σ . Then it follows that

$$\begin{aligned} \mathbb{E} [r^\top x] &= \mu^\top x \\ \text{Var} [r^\top x] &= x^\top \Sigma x \end{aligned}$$

In words, the expected return, which is the expectation of the return $r^\top x$, is given by $\mu^\top x$. Moreover, the *risk* of my portfolio, which is often defined by the variance of the return $r^\top x$, is given by $x^\top \Sigma x$. By definition, the covariance matrix is positive semidefinite.

We want to find a portfolio that maximizes the expected return while guaranteeing a low risk. Then we consider

$$\begin{aligned} &\text{maximize} && \mu^\top x - \gamma x^\top \Sigma x \\ &\text{subject to} && 1^\top x = 1, \\ &&& x \in C' \end{aligned}$$

where $\gamma > 0$ is the risk aversion parameter. When $C' = \mathbb{R}_+^d$, we take long positions only. When $C' = \{x \in \mathbb{R}^d : \|x\|_1 \leq B\}$, then we allow short positions but there is a leverage limit.

4.2 Uncertainty quantification

Suppose we have chosen a portfolio x . To avoid high risk portfolios, can we measure the worst-case variance of the given portfolio? In practice, the covariance matrix Σ is estimated through data, so it is subject to errors. Given a magnitude ϵ of potential errors, what is the highest risk of the portfolio?

$$\begin{aligned} & \text{maximize} && x^\top (\Sigma + S)x \\ & \text{subject to} && S \succeq 0, \\ & && \|S\|_{\text{nuc}} \leq \epsilon \end{aligned}$$

where $\|S\|_{\text{nuc}}$ denotes the *nuclear norm* of S , defined as the sum of all eigenvalues of S . Is this problem convex?

4.3 Support vector machine

Given n data $(x_1, y_1), \dots, (x_n, y_n)$ where $y_i \in \{-1, 1\}$ are labels, we want to find a separating hyperplane

$$w^\top x = b$$

to classify data with $+1$ and data with -1 . The goal is to find a separating hyperplane $w^\top x = b$ with the “gap” ($1/\|w\|_2$) being maximized. Then the problem can be formulated as

$$\begin{aligned} & \text{minimize} && \|w\|_2 \\ & \text{subject to} && y_i(w^\top x_i - b) \geq 0, \quad i = 1, \dots, n. \end{aligned}$$

If this problem is feasible, then $x \rightarrow \text{sign}(w^\top x_i - b)$ is a valid classifier for the data set.

What if the data set is not entirely separable? What if no hyperplane separates the data without an error? In such cases, we force separation via a penalty term, instead of imposing hard constraints. The number of misclassifications can be used as penalty. Namely,

$$\sum_{i=1}^n 1(y_i \neq \text{sign}(w^\top x_i - b)).$$

However, this is not convex. Instead, we apply the *hinge loss*¹, which is an upper bound on the number of misclassifications, given by

$$\sum_{i=1}^n \max\{0, 1 - y_i(w^\top x_i - b)\}.$$

Then we solve

$$\min_{w,b} \lambda \|w\|_2^2 + \frac{1}{n} \sum_{i=1}^n \max\{0, 1 - y_i(w^\top x_i - b)\}$$

where λ determines the trade-off between the margin size and the penalty.

¹Here, $\max\{0, a\}$ is called the hinge function.